

and it is said to extend near to the southern limits of the Union. Michaux, in his account of the forest trees, states, that it is particularly abundant through the whole range of Allegany mountains, upon the borders and near the sources of rivers. It gradually diminishes however on both sides as these rivers approach to the sea, or to their confluence with the great western streams.

The botanical character of the genus consists in a *five parted calyx, a hypocrateriform corolla, containing ten depressions in its border, in which the anthers are lodged; a capsule five celled.*

The specific character is, that the *leaves are scattered, petioled, oval and smooth; the corymbs terminal, viscid and pubescent.*

Class *Decandria*, order *Monogynia*. Natural orders *Bicornes*, Linn. *Rhododendra*, Juss.

The height of the *Kalmia latifolia* is generally that of a shrub, sometimes however attaining to the altitude of a small tree. Its leaves are evergreen, coriaceous, very smooth, with the under side somewhat palest. Their form is oval, acute and entire; their insertion by scattered petioles, on the sides and extremities of the branches. The flowers vary from white to red; they grow in terminal corymbs, simple or compound with opposite branches, and made up of slender peduncles. These

are invested with a glutinous pubescence, and supported at base by ovate, acuminate bractes. The calyx is small, five parted, persistent, with oval, acute segments. The corolla is monopetalous, with a cylindrical tube, a spreading disc, and an erect, five cleft margin. At the circumference of the disc on the inside, are ten depressions or pits, accompanied with corresponding prominences on the outside. In these depressions the anthers are found lodged at the time when the flower expands. The stamens originate from the base of the corolla, and bend outwardly, so as to lodge their anthers in the cells of the corolla. From this confinement they liberate themselves during the period of flowering and strike against the sides of the stigma. The germ is roundish, the style longer than the corolla and declined, the stigma obtuse. Capsule roundish, depressed, five celled and five valved, with numerous small seeds.

I have examined chemically the leaves of the *Kalmia*, gathered at the time the shrub was in fruit. The following constituent principles were found to exist in them.

1. Vegetable mucus. This exists in large quantities, and is dissolved in water both by infusion and decoction, rendering it extremely mucilaginous or ropy. When alcohol is added to this so-